



GRAMIN KRISHI MAUSAM SEWA
Agromet Advisory Bulletin for Sub Tropical Conditions of Jammu Region
Issued Jointly by Sher-e-Kashmir University of Agricultural Sciences & Technology Jammu and
India Meteorological Department
Agromet Research Centre, SKUAST-Jammu, Chatha-180009
email:jmu.agrimet@gmail.com



No. AUJ/Agromet/GKMS/F-1/18-19/227	Bulletin No: 39/2018-19	Dated: 14-08-2018
Phone:0191-2263891	Website: www.skuast.org	Fax: 0191-2263891

District:Samba

Weather summary of previous 5 days (09th to, 13th August 2018)	Range of Max. Temperature (⁰C)	NA				
	Range of Min. Temperature (⁰C)	NA				
	Range of Relative Humidity (%)	NA				
	Rainfall (mm)	12.8				
Ensembled forecast for next 5 days (0830 hrs, 15th to 19th Aug, 2018)	Date	15/08/18	16/08/18	17/08/18	18/08/18	19/08/18
	Rainfall (mm)	25	30	15	5	5
	Max. temperature (⁰C)	30	30	30	33	33
	Min. temperature (⁰C)	22	22	22	24	24
	Total cloud cover (Okta)	7	7	7	7	7
	RH at 08:30 Hrs (%)	95	95	95	95	95
	RH at 17:30 Hrs (%)	80	80	80	80	80
	Wind speed (km/hr.)	1	1	1	1	1
	Wind direction (degrees)	120	95	150	210	95

WEATHER FORECAST BASED AGROMET ADVISORIES W.E.F. 15th to 19th Aug, 2018

Crop	Stage	Weather based Agro- advisories
Maize	Early vegetative/ knee high	Keeping in view the rainfall in past days and next 5 days, farmers are advised to make arrangement to drain out excess water from the fields. Farmers may apply fertilizer as well as chemical plant protection measures on 19 th August after receipt of rain.
Kharif Pulses (moong/mash)	Germination/ early vegetative/	Keeping in view the rainfall in past days and next 5 days, farmers are advised to make arrangement to drain out excess water from the fields. Farmers may apply fertilizer as well as chemical plant protection measures on 19 th August after receipt of rain.
Paddy	Establishment/ Vegetative	Keeping in view the rainfall in past days and next 5 days, postpone irrigation due to chances of rain. Make arrangement to drain out excess water from the fields. Farmers may apply 2 nd dose of nitrogenous fertilizer as top dressing in normal/early transplanted rice crop on 19 th August after receipt of rain.
Kharif mixed fodder	Early vegetative	Keeping in view the rainfall in past days and next 5 days, farmers are advised to make arrangement to drain out excess water from the fields. Postpone application of fertilizer as well as chemical plant protection measures till rainfall of 19 th August.
Vegetables: Okra	Harvesting	Keeping in view the rainfall in past days and next 5 days, farmers are advised to make arrangement to drain out excess water from the fields. Farmers may apply fertilizer as well as chemical plant protection measures on 19 th August after receipt of rain.
Bitter gourd/bottle gourd	Harvesting	
Turmeric	Vegetative	
Rabi season crop	Nursery under protection	
Floriculture: Chrysanthemum	Cutting preparation	Keeping in view the rainfall in past days and next 5 days, farmers are advised to make arrangement to drain out excess water from the fields. Farmers may apply fertilizer as well as chemical plant protection measures on 19 th August after receipt of rain.
Marigold	Sowing	
Horticulture Mango	Bearing	Make arrangement to drain out excess water from the fields. Farmers may apply fertilizer as well as chemical plant protection measures on 19 th August after receipt of rain.



GRAMIN KRISHI MAUSAM SEWA
Agromet Advisory Bulletin for Sub Tropical Conditions of Jammu Region
Issued Jointly by Sher-e-Kashmir University of Agricultural Sciences & Technology Jammu and
India Meteorological Department
Agromet Research Centre, SKUAST-Jammu, Chatha-180009
email:jmu.agrimet@gmail.com



Mushroom Milky mushroom		Prepare casing soil, if spawn run is complete. Maintain temperature at 20-25°C
Oyster mushroom	Fruit initiation	Pinch 2 mm diameter holes in full run spawn bags of oyster mushroom.
Sericulture Mulberry	Tree plantation	Water logging in newly transplanted mulberry saplings should be avoided by digging proper exit channels.
Cocoon crop	Harvested cocoons	Cocoon harvested during April month (if not sold) should be loosely packed in muslin cloth bags. The bags should be kept in wire mesh covered racks to avoid rodent and fungal attack till sale of the crop.
Apiculture		Migrate bee colonies to safer areas having bee flora. Provide sugar feeding/artificial diet. Protect bee hive from wax moth and ants. Unite weak bee colonies. Protect colonies from rain.
Poultry		Water supply should be adequate and add vitamin C in water. Proper ventilation must be maintained at poultry house. Fumigate poultry huts. Maintain hygiene. Birds should not come in direct contact with rain. Use fan in poultry houses due to high humidity.
Live stock: Dairy animals/ Sheep & Goat		Increase protein content in animal ration by 2%. Mineral mixture should be supplemented in diet. Give vaccination against FMD. Provide clean drinking water to animals.
Fisheries Culture fisheries	Yearlings/ Fingerlings/fry	Apply the screen/netting cloth on the outlets/overflows so that fish stock may not escape out during excessive rains.

Postpone application of fertilizer as well as chemical plant protection measures till weather gets clear. Drain out excess water from fields.

Dr Mahender Singh
I/C Agrometeorology

Dr Veena Sharma
Technical Officer